

Combinatorial and topological phase structure of non-perturbative n -dimensional quantum gravity

M.Carfora*, M.Martellini**, A.Marzuoli*

Dipartimento di Fisica Nucleare e Teorica dell'Università di Pavia
Via Bassi 6, I-27100 Pavia, Italy*

Istituto Nazionale di Fisica Nucleare, Sezione di Roma I, Italy,
and Dipartimento di Fisica dell' Università di Milano

Via Celoria 16, I-20133 Milano, Italy (permanent address) **
Istituto Nazionale di Fisica Nucleare, Sezione di Pavia, Italy*, **

February 7, 2008

Abstract

We provide a non-perturbative geometrical characterization of the partition function of n -dimensional quantum gravity based on a coarse classification of riemannian geometries. We show that, under natural geometrical constraints, the theory admits a continuum limit with a non-trivial phase structure parametrized by the homotopy types of the class of manifolds considered. The results obtained qualitatively coincide, when specialized to dimension two, with those of two-dimensional quantum gravity models based on random triangulations of surfaces.

1 Introduction

In this paper we present results which extend to any dimension $n \geq 2$ some of the non-perturbative aspects of two-dimensional quantum gravity theories based on random triangulations of surfaces^{1, 2, 3}.

This extension relies on previous work of two of us on functional integration on the space of riemannian structures^{4, 5}, and it is made possible by exploiting the properties of the space of bounded geometries introduced by M. Gromov⁶. In particular, we shall make use of the infinite dimensional set, $\mathcal{R}(n, r, D, V)$, of all compact riemannian manifolds of dimension n (not necessarily of the same underlying topology), with sectional curvature bounded below by $-(n-1)r$, diameter bounded above by D and volume bounded below by V (r being any real number, and D, V any positive real numbers)⁶. $\mathcal{R}(n, r, D, V)$ is an infinite-dimensional space which can be endowed with a natural topology most suitable for handling questions of convergence of random triangulations on riemannian manifolds. A property, this latter, connected to the observation that for any manifold M in such a class it is possible to introduce coverings by geodesic balls whose combinatorial patterns yields simplicial approximations which provide a coarse classification of the riemannian structures occurring in $\mathcal{R}(n, r, D, V)$ ⁷.

In what follows we need few details on such coverings, thus, let $L(m) \equiv 1/m$, denote a constant (later on to be interpreted as a cut-off related to the spacing of a triangulation). For any given $L(m)$, and any given manifold M in $\mathcal{R}(n, r, D, V)$, we can always introduce on M geodesic balls coverings in such a way that a finite collection of open geodesic balls of radius $2L(m)$ covers M , while the corresponding balls of radius $L(m)$ are disjoint, (this particular covering is often called a minimal $L(m)$ -net for M). Two combinatorial invariants characterize such construction. The first is the *filling function* $N_{(m)}(M)$ of the covering, *i.e.*, the function which associates with M the maximum number of geodesic balls realizing a minimal $L(m)$ -net on M .

The second invariant is the (first) *intersection pattern* of the covering, $\gamma_{(m)}(M)$, defined by the collection of pairs of geodesic balls of radius $2L(m)$ having a non empty pairwise intersection. These two invariants characterize the 1-skeleton, $\Gamma_{(m)}(M)$, of the geodesic balls covering. Similarly, upon considering the higher order intersection patterns, one ends up in defining the two-skeleton $K_2(M)$, and eventually the *nerve* of the covering of the manifold M . If m is sufficiently large this nerve gives rise to a polytope which can be assimilated to a symplcial approximation of the manifold M . In particular, the 1-skeleton is just the vertex-edge structure of this approximation.

Any two manifolds M_1 and M_2 in $\mathcal{R}(n, r, D, V)$ endowed with minimal $L(m)$ -net are considered equivalent if and only if they have the same filling functions and the same intersection patterns up to combinatorial isomorphisms. Such relation partitions $\mathcal{R}(n, r, D, V)$ into disjoint equivalence classes whose number can be shown to be *finite*,

function of the parameters m, n, r, D, V ⁷. Each equivalence class of manifolds is characterized by the abstract (unlabelled) graph $\Gamma_{(m)}$ defined by the 1-skeleton of the $L(m)$ -covering. The order of any such graph (*i.e.*, the number of vertices) is provided by the filling function $N_{(m)}$, while the structure of the edge set of $\Gamma_{(m)}$ is defined by the intersection pattern $\gamma_{(m)}(M)$. It is important to remark that on $\mathcal{R}(n, r, D, V)$ either the filling function or the intersection pattern cannot be arbitrary. The former is always bounded above for each given m , (for instance, if $r < 0$ an upper bound is provided by $D^n m^n$), and the best filling of a riemannian manifolds with geodesic balls of radius $1/m$ is realized on (portions of) spaces of constant curvature ⁶. The latter is similarly controlled through the geometry of the manifold to the effect that the *average degree*, $d(\Gamma)$, of the graph $\Gamma_{(m)}$, (*i.e.*, the average number of edges incident on a vertex of the graph), approaches a constant value as the radius of the balls defining the covering tend to zero, (*i.e.*, as $m \rightarrow \infty$). Such constant is independent from m , and can be estimated in terms of the parameters n, r, D , and V ⁷.

An intuitive idea of the natural topology of the Gromov space of bounded geometries can be grasped by noticing that two riemannian manifolds in $\mathcal{R}(n, r, D, V)$ get closer and closer in such topology if we can introduce on them finer and finer minimal $L(m)$ -nets of geodesic balls with the same filling function and the same intersection pattern, *i.e.*, if the symplcial approximations induced by geodesic balls coverings have a similar vertex-edge structure.

Within this geometrical framework, let us consider the following graph-theoretical statistical sum

$$\Xi(m, z) = \sum_{\Gamma} z^{N_{(m)}(\Gamma)} \exp[\beta \Phi_{(m)}(\Gamma)] \quad (1)$$

where $z \equiv \exp(-c)$, and β are constants, and where the (finite) sum is over all inequivalent 1- skeletons $\Gamma_{(m)}$, with order $N_{(m)}(\Gamma)$ and size $\Phi_{(m)}(\Gamma)$, realized by the possible $L(m)$ -geodesic balls covering over manifolds in $\mathcal{R}(n, r, D, V)$, (the size $\Phi_{(m)}(\Gamma)$ is the number of edges in the graph $\Gamma_{(m)}$). Notice that either the filling function $N_{(m)}$ or the size $\Phi_{(m)}(\Gamma)$, besides having a natural combinatorial meaning for the manifolds in $\mathcal{R}(n, r, D, V)$, can be directly related to riemannian invariants. A standard expansion for the riemannian volume of small geodesic balls yields $N_{(m)}(M) \simeq B(n, m)\{1 + [m^{-2}/6(n+2)]R + o(m^{-2})\}$, where $B(n, m)$ is a constant depending on the volume of M , R is the scalar curvature evaluated at a suitable point and $o(m^{-2})$ stands for higher order terms which are universal polynomials in the curvature tensor and its covariant derivatives. A similar expansion holds true also for $\Phi_{(m)}(\Gamma)$. In particular, from the properties of geodesic balls coverings it follows that the size of the graph Γ grows with m as $N_{(m)}$ does, and for m sufficiently large one can write $\Phi_{(m)}(\Gamma(M)) \simeq B_{M,R} N_{(m)}(\Gamma(M))$, where $B_{M,R}$ is a constant, independent from m , related to the curvature of M . Collecting such results, we can heuristically rewrite

$\Xi(m, z)$, to leading order in m , as

$$\Xi(m, z) \simeq \sum_{\Gamma} \exp[m^n(-c \cdot Volume + \beta \cdot Curvature)]$$

Since the above summation over all graphs Γ is a sum over all finite equivalence classes of manifolds in $\mathcal{R}(n, r, D, V)$ parametrized by their one-skeletons, we can regard $\Xi(m, z)$ as a discrete version of a sum over manifolds in $\mathcal{R}(n, r, D, V)$ with a combinatorial weight related to volume and curvature.

2 Geodesic balls coverings and lattice gas statistical mechanics

From a thermodynamical point of view, $\Xi(m, z)$ has the structure of a grand-partition function computed for a lattice gas with negative pair interactions evaluated at inverse temperature β and for a fugacity z .

In order to discuss this statistical equivalence more in details we explicitly identify the geodesic balls of a minimal $L(m)$ -net on a manifold M of bounded geometry with the vertices $\{p_1^{(0)}, \dots, p_N^{(0)}\}$, of an abstract graph $\Gamma(M)$. Such $\Gamma(M)$ is defined by connecting the $\{p_i^{(0)}\}$ among themselves with undirected edges $p_{ij}^{(1)} = \{p_i^{(0)}, p_j^{(0)}\}$ if and only if the geodesic balls labelled $p_i^{(0)}$ and $p_j^{(0)}$ have a non-empty intersection when we double their radius. Any two such graph are considered equivalent if, up to the labelling $\{p_i^{(0)}\}$ of the balls, they have the same vertex-edge scheme, *viz.* we are considering unlabelled graphs associated with the possible one-skeletons of geodesic balls coverings of manifolds of bounded geometries. For any given m , any such graph can be topologically imbedded in a larger regular, (*i.e.*, whose vertices have all the same degree), graph $\Omega_{(m)}$, (Ω for short), having as its connected subgraphs all possible 1- skeletons $\Gamma_{(m)}$ which are realized by $L(m)$ -geodesic balls coverings as M varies in $\mathcal{R}(n, r, D, V)$. One may think of such Ω as the configuration space for the finite set of possible 1-skeletons that can be realized by minimal nets on the manifolds in $\mathcal{R}(n, r, D, V)$.

For a given m , a graph realizing Ω can be constructively defined first by labelling the graphs $\Gamma_{(m)}$ and order them by inclusion, *i.e.*, $\Gamma_{(m)}^{(i)} \subset \Gamma_{(m)}^{(j)}$ if $\Gamma_{(m)}^{(i)}$ is a subgraph of $\Gamma_{(m)}^{(j)}$, and then by considering the union $H_{(m)}$ over the *prime* graphs $\Gamma_{(m)}^{(k)}$, (*i.e.*, over those graphs which do not appear as subgraphs of other $\Gamma_{(m)}$):

$$H_{(m)} \equiv \cup_j^* \Gamma_{(m)}^{(j)} \quad (2)$$

Let $d(\Omega_{(m)})$ the maximum degree of the vertices of H , (or which is the same, the maximum degree of the vertices among those occurring in the various $\Gamma_{(m)}^{(j)}$). Notice that, for every m , we have $d(\Omega) \leq C_{n,r,D}$, where the constant $C_{n,r,D}$ is the upper bound to the average degree of the graphs $\Gamma_{(m)}$ associated with the $L(m)$ -geodesic balls realized in $\mathcal{R}(n, r, D, V)$.

The existence of the graph $\Omega_{(m)}$ is assured by a theorem of P.Erdos and P.Kelly⁸, and in order to construct it, we proceed as follows⁴. First add to $H_{(m)}$ a set I of $b_{(m)}$ isolated points, (the number of which, as indicated, is a function of the given m), a new graph is formed from $H_{(m)}$ and I by adding edges between pairs of points in I and $H_{(m)}$, (notice that no edges are added between vertices in $H_{(m)}$). The strategy is to give rise, in this way, to a new graph which is regular of degree $d(\Omega_{(m)})$ while keeping the number $b_{(m)}$ of vertices to be added to $H_{(m)}$ as small as possible. As expected, the number $b_{(m)}$ of added vertices depends only on the degree sequence of the graph $H_{(m)}$. In particular if d_i denote the various degrees at the vertices of $H_{(m)}$, then $b_{(m)}$ is the least integer satisfying: (i) $b_{(m)}d(\Omega_{(m)}) \geq \sum_i (d(\Omega) - d_i)$, (ii) $b_{(m)}^2 - (d(\Omega) + 1)b_{(m)} + \sum_i (d(\Omega) - d_i) \geq 0$, (iii) $b_{(m)} \geq \max(d(\Omega) - d_i)$, and (iv) $(b_{(m)} + h_{(m)})d(\Omega)$ is even, where $h_{(m)}$ is the order of the union graph $H_{(m)}$ ⁸.

It should be noted that in the large m limit, $\Omega_{(m)}$ is a regular graph whose degree, $d(\Omega_{(m)})$, is independent from m , being bounded above in terms of n, r, D , by $(C_{n,r,D})^{n-1}$.

With these preliminary remarks we can work out the correspondence between minimal geodesic balls coverings of manifolds M in $\mathcal{R}(n, r, D, V)$ and the statistical system described by a lattice gas on $\Omega_{(m)}$ by identifying the collection of geodesic balls, providing the dense packing of M , with a gas of indistinguishable particles. Each particle can occupy at most one site of $\Omega_{(m)}$, (the geodesic balls of radius $L(m) = 1/m$ are disjoint), and the configuration of occupied sites corresponding to the net of balls covering M is defined by the vertices of the graph $\Gamma(M) \subset \Omega_{(m)}$.

The interaction energy between two occupied sites is supposed to be different from zero only if the sites in question are connected by an edge $\{p_i^{(0)}, p_j^{(0)}\}$ of $\Gamma(M)$, namely if the points of the minimal $L(m)$ -net corresponding to the lattice sites in question are in the intersection pattern of the manifold according to the definition recalled above, (the geodesic balls $p_i^{(0)}$ and $p_j^{(0)}$ have a non-empty intersection when their radius is doubled). More in general, $p_i^{(0)}$ and $p_j^{(0)}$ will be said to be neighbors if $(p_i^{(0)}, p_j^{(0)})$ is an edge of the graph $\Omega_{(m)}$. Obviously, $\Omega_{(m)}$ will contain as possible configurations of occupied sites, (*i.e.*, as possible subgraphs), not only those graphs $\Gamma(M)$ representing one-skeleton of manifolds in $\mathcal{R}(n, r, D, V)$, but also graphs associated to the $b(m)$ added vertices (and to the corresponding edges), needed in order to construct $\Omega_{(m)}$. In any case, given a configuration of occupied sites in $\Omega_{(m)}$ represented by a graph Γ with $N_{(m)}(\Gamma)$ vertices, the total interaction energy corresponding to such configuration is given by

$$E(\Gamma) = -e_0 \Phi_{(m)}(\Gamma) \quad (3)$$

where e_0 is a constant (which provide the scale of energy, and which here and henceforth we set equal to one) and $\Phi_{(m)}$ is the number of edges in the graph Γ , *viz.*, when dealing with one-skeleton graphs, the total number of pairs (i, j) belonging to the intersection pattern of the minimal $L(m)$ -net $\{p_1^{(0)}, \dots, p_N^{(0)}\}$. The corresponding Boltzmann weight

(as a lattice gas) is then

$$\exp[-\beta E(\Gamma)] = \exp[\beta \Phi_{(m)}(\Gamma)] \quad (4)$$

Roughly speaking, according to the correspondence just established the configurations of the lattice gas considered are labelled by random graphs representing the possible 1-skeletons $\Gamma_{(m)}$ of manifolds with bounded geometry, and, for a sufficiently large m , we get a grand-ensemble of such configurations whose thermodynamical parameters keep track of the average volume and curvature of the manifolds in $\mathcal{R}(n, r, D, V)$.

It must be stressed that here we are dealing with a statistical system not evolving on a regular lattice, for, it shows up more complex interactions which correspond to the non-trivial topology of the manifolds underlying the coverings. For manifolds in $\mathcal{R}(n, r, D, V)$, the boundedness properties for the filling function and the cardinality of the intersection pattern imply that such interactions are localized and do not yield for a catastrophic behavior when going to the thermodynamic limit. From a more technical side, one proves the existence of such limit by exploiting the compactness properties of $\mathcal{R}(n, r, D, V)$. These properties easily allow to show that $\lim_{m \rightarrow \infty} \Xi(m, z)$ exists even if in general this limit is not unique. Such non-uniqueness corresponds to the possible onset of phase transitions in the system. To be more specific, let us take advantage of the introduction of the graph $\Omega_{(m)}$ in order to rewrite $\Xi(m, z)$ as the polynomial

$$\Xi(m, z) = \sum_{\Gamma} \hat{z}^{N_{(m)}(\Gamma)} \prod_{p \in \Gamma} \prod_{q \in \Omega \setminus \Gamma} \exp\left[-\frac{1}{2} \beta A_{\Omega \setminus \Gamma}(p, q)\right] \quad (5)$$

where $A_{\Omega \setminus \Gamma}(p, q)$ is the adjacency matrix of the graph $\Omega \setminus \Gamma$, (*i.e.*, the matrix whose entries are 1 if the vertices p and q of the graph in question are connected by an edge, zero otherwise; with a slight abuse of notation, we have denoted by $\Omega \setminus \Gamma$ the graph in Ω obtained by removing all the edges, but not the vertices, belonging to Γ). We have also introduced a *normalized fugacity* $\hat{z}$ according to

$$\hat{z} \equiv z \exp\left[\frac{1}{2} \beta d(\Omega)\right] \quad (6)$$

(Notice that in general $\hat{z}$ depends from the given m through the expression of the degree $d(\Omega_{(m)})$, however, for m sufficiently large this dependence will eventually disappear). We wish to stress again that the sum (5) is *restricted* only to those graphs $\Gamma_{(m)}$ which are $L(m)$ -geodesic balls one-skeletons for manifolds in $\mathcal{R}(n, r, D, V)$, and it is *not extended* to all possible subgraphs of $\Omega_{(m)}$. The *unrestricted sum*, $\Xi(m, \hat{z})^*$, is given by

$$\Xi(m, \hat{z})^* = \sum_{\forall \Gamma \subset \Omega} \hat{z}^{N_{(m)}(\Gamma)} \prod_{p \in \Gamma} \prod_{q \in \Omega \setminus \Gamma} \exp\left[-\frac{1}{2} \beta A_{\Omega \setminus \Gamma}(p, q)\right] \quad (7)$$

and we can formally write

$$\Xi(m, \hat{z})^* = \Xi(m, \hat{z}) + \sum_{\Omega \setminus f(\mathcal{R}(n, r, D, V))} \hat{z}^{N_{(m)}(\Gamma)} \prod_{p \in \Gamma} \prod_{q \in \Omega \setminus \Gamma} \exp[-\frac{1}{2} \beta A_{\Omega \setminus \Gamma}(p, q)] \quad (8)$$

where the sum over $\Omega \setminus f(\mathcal{R}(n, r, D, V))$ indicates summation over all graphs in $\Omega_{(m)}$ which are not $L(m)$ -one-skeletons for manifolds in $\mathcal{R}(n, r, D, V)$.

Notice that in the construction of $\Omega_{(m)}$ out of the one-skeleton graphs $\Gamma_{(m)}(M)$ with the Erdos-Kelly algorithm, (see (2)), no edges are introduced between the graphs $\Gamma_{(m)}(M)$, new edges may occur only between the $b_{(m)}$ added vertices and between these latter vertices and the graphs $\Gamma_{(m)}(M)$. From such remarks it follows that the graphs in $\Omega_{(m)} \setminus f(\mathcal{R}(n, r, D, V))$ are defined by the $b_{(m)}$ added vertices and by the possible edges between them and the graphs $\Gamma_{(m)}$.

The relation (8) between the unrestricted, Ξ^* , and the restricted statistical sum Ξ , is useful for suggesting the boundary conditions which uncover the non-trivial phase structure of $\lim_{m \rightarrow \infty} \Xi(m, \hat{z})$.

In order to proceed in this direction, let us notice that the unrestricted statistical sum (7), $\Xi(m, \hat{z})^*$, takes on the classical polynomial Lee-Yang structure⁹. It follows, according to the Lee-Yang circle theorem⁹, that the zeroes of $\Xi(m, \hat{z})^*$, thought of as a function of the complex variable $\hat{z}$, all lie, for each given value of m , on the circle $\{\hat{z}: |\hat{z}| = 1\}$ in the plane of the complexified fugacity $\hat{z}$.

3 Critical behavior

The Lee-Yang type result quoted above relates the presence or absence of a phase transition to the analyticity properties of the free energy associated with the grand-partition function Ξ^* . A rather delicate analysis⁴, shows that it is possible to carry out the thermodynamic limit on Ξ^* , with the boundary condition of keeping empty the $b_{(m)}$ added sites. (Geometrically speaking, this boundary condition favors the sampling of subgraphs of $\Omega_{(m)} \uparrow \Omega_\infty$ which represents geodesic balls coverings of manifolds in $\mathcal{R}(n, r, D, V)$). Then, on applying a variant of the Pirogov- Sinai theorem⁹ on the occurrence of phase transitions in lattice gases with attractive pair interactions, one can show that at sufficiently low temperature the statistical system described by Ξ undergoes a phase transition, in the large m limit, as $\hat{z} \rightarrow 1^-$.

The interpretation in geometrical terms of the critical behavior of $\Xi(m, \hat{z})$ as $m \rightarrow \infty$ and $\hat{z} \rightarrow 1^-$ relies on a deep result on the structure of $\mathcal{R}(n, r, D, V)$. This result⁷ states that, for m sufficiently large, manifolds in $\mathcal{R}(n, r, D, V)$ with the same 1-skeleton $\Gamma_{(m)}$ are homotopically equivalent, and the number of different homotopy-types of manifolds realized in $\mathcal{R}(n, r, D, V)$ is finite. In particular, if m_0 is the smallest m such that geodesic balls of radius smaller than $2L(m_0)$ are contractible, then $\mathcal{R}(n, r, D, V)$ contains less than $[\hat{N}_{(m_0)}]^4$ distinct homotopy types, where $\hat{N}_{(m)}$ denotes an upper bound to the filling

function over $\mathcal{R}(n, r, D, V)$. In connection with this homotopy finiteness theorem, it can be explicitly shown⁴ that the continuum limit of the sequence of measures associated with $\Xi(m, z)$ yields for at least two limit distributions for sufficiently low temperatures. Such distributions are parametrized by the fundamental group of the manifolds sampled in $\mathcal{R}(n, r, D, V)$ by Ξ .

It is possible to provide a less formal interpretation of the critical behavior of Ξ which, even if less rigorous than the one just sketched, has the advantage of a more direct connection with the two-dimensional theory. To this end let us introduce the function $B_\lambda(\Gamma_{(m)}, \pi(M))$ which counts the number of combinatorially inequivalent 1-skeletons $\Gamma_{(m)}$ with λ vertices and with given average degree d_λ , which can be drawn on a manifold M in the homotopy class $\{\pi(M)\}$. This function depends only on the fundamental group $\pi_1(M)$ of M . Its asymptotic behavior, as $\lambda \rightarrow \infty$, is well known for surfaces of given genus¹⁰, and it is one of the main ingredient for understanding the double-scaling limit in two-dimensional quantum gravity models. A qualitatively similar behavior can be obtained here too for B_λ , by rewriting $\Xi(m, z)$ as $\sum_\lambda \hat{z}^\lambda a_\lambda$ with $a_\lambda \equiv \sum_{\{\pi(M)\}} \sum_{d_\lambda} B_\lambda(\Gamma_{(m)}, \pi(M)) \exp[1/2\beta(d_\lambda - d(\Omega))\lambda]$.

Notice that the summation over the average degrees, $\sum_{d_\lambda}$, extends, independently of λ , over a finite number of terms since we have the uniform bound $d_\lambda \leq C_{n,r,D}$, which does not depend on m , (the possibility of the above rewriting readily follow from (1) by expressing the sum over the 1- skeleton $\Gamma_{(m)}$ as the *finite* sum over the distinct homotopy types $\{\pi(M)\}$ sampled in $\mathcal{R}(n, r, D, V)$). By Lee-Yang theorem, the power series $\Xi(\hat{z}) \equiv \lim_{m \rightarrow \infty} \sum_\lambda \hat{z}^\lambda a_\lambda$ has radius of convergence one; *i.e.*, $\limsup_{\lambda \rightarrow \infty} (a_\lambda)^{1/\lambda} = 1$, which yields, for each homotopy type,

$$\limsup_{\lambda \rightarrow \infty} \frac{\ln B_\lambda(\Gamma_{(m)})}{\lambda} = \limsup_{m \rightarrow \infty} \frac{1}{2} \beta [d(\Omega_{(m)}) - d_\lambda] \equiv \ln \Lambda_c$$

(for the sake of simplicity, here we are tacitly assuming that for a given λ , just one average degree d_λ is dominating in a_λ . The argument can be extended to the situation where distinct degrees do contribute. In the two-dimensional theory, this latter circumstance corresponds to surfaces discretized with different percentages of n -gones, say triangles, squares, and pentagons).

The above relation implies that asymptotically in λ , B_λ behaves as $(\Lambda_c)^\lambda \exp o(\lambda)$. The geometrical meaning of the terms $o(\lambda)$ in this asymptotics follows by noticing that $\Gamma_{(m)}$ is a graph drawn on a two-dimensional complex $K_2(M)$, (the two-skeleton of M), whose fundamental group is isomorphic to $\pi_1(M)$. This implies that the number of edges of $\Gamma_{(m)}(M)$ is given by $\Phi_{(m)}(\Gamma(M)) = \lambda + h[\pi_1(M)] + \rho(\Gamma) - 1$, where $h[\pi_1(M)]$ is the number of non-trivial generators of $\pi_1(M)$, and where $\rho(\Gamma)$ is the number of relations in $\pi_1(M)$ obtained by pasting faces onto the graph so as to get $K_2(M)$. [To be more precise, the inclusion $\Gamma_{(m)} \rightarrow K_2$ induces an epimorphism between the free fundamental group of the graph $\Gamma_{(m)}$ and the fundamental group of K_2 . This implies that this latter fundamental group, which is isomorphic to the fundamental group of the manifold M , has

a presentation with $1 + \Phi_{(m)}(\Gamma) - N_{(m)}(\Gamma)$ generators and $\rho(\Gamma)$ relations. It can be shown that $h[\pi_1(M)]$ is the (finite) number of geodesic loops generators, (with given basepoint), non homotopic to zero, representing the inequivalent homotopy classes of $\pi_1(M)$.

From the properties of the geodesic balls coverings it follows that $\lim_{\lambda \rightarrow \infty} \Phi_{(m)}/\lambda$ exists and it is finite. Thus, for large λ , and for any manifold M in $\mathcal{R}(n, r, D, V)$, we can write $\rho(\Gamma(M)) = \alpha_h(M)\lambda + \sigma_h(M)$ where we have introduced the constants α_h and σ_h which in general depend on the number of non trivial generators h of $\pi_1(M)$. According to these remarks we can eventually write the number of edges of $\Gamma(M)$ in terms of the non-trivial generators of π_1 as

$$\Phi_{(m)}(\Gamma(M)) = (1 + \alpha_h(M))\lambda + h[\pi_1(M)] + \sigma_h(M) - 1 \quad (9)$$

Such an expression shows that with each graph Γ , representing possible one-skeletons of geodesic balls coverings, we may formally associate a graph $\hat{\Gamma}$ with the same number of edges but with $\hat{\lambda} \equiv (1 + \alpha_h(M))\lambda$ vertices. In particular, it follows that the counting function $B_\lambda(\Gamma_{(m)})$ is proportional to the number of unlabelled graphs with $\hat{\lambda}$ vertices and $\Phi_{(m)}$ edges.

In order to count such graphs we can use Pólya's enumeration theorem ¹¹. A direct computation allows to show that the following estimate holds true

$$C_1\{(C_2)^\lambda \lambda^{h[\pi_1(M)] + \sigma_h - 5/2}\} \leq B_\lambda \leq C_3\{(C_4)^\lambda \lambda^{h[\pi_1(M)] + \sigma_h - 3/2} + O(\lambda^{3/2} \exp[\lambda^2])\}$$

where C_1, C_2, C_3, C_4 denote suitable constants. A comparison of this estimate with the asymptotics for B_λ , derived from Lee-Yang theorem, allows to show that the term $\exp[o(\lambda)]$ in such asymptotics is given by $\lambda^{h[\pi_1(M)] + \sigma_h - \xi}$, where ξ is a parameter with $3/2 \leq \xi \leq 5/2$.

Since we are counting inequivalent graphs which can be drawn on a two-dimensional object, (the two-skeleton K_2), the asymptotics for B_λ , as derived above, is, as expected, consistent with the asymptotics that could have been obtained by graphs counting matrix-models techniques. More in particular, if we set, (as we do henceforth, for simplicity), $\xi = 2 - \gamma_0$ and $\sigma_h(M) = -(1/2)\gamma_0 h[\pi_1(M)] - 1$ where γ_0 is a parameter with $-(1/2) \leq \gamma_0 \leq (7/2)$, then $B_\lambda \sim \rho_h(\Lambda_c)^\lambda \lambda^{(1-h/2)(\gamma_0-2)-1}$, (where ρ_h is a suitable constant), which is the usual asymptotics for the number of unlabelled graphs (with λ vertices) that can be drawn on a two-dimensional surface of genus $(1/2)h[\pi_1(M)]$.

According to such asymptotic behavior for B_λ , a Tauberian theorem, (actually an Abelian regularity results), yields for $\Xi(\hat{z})$, as $\hat{z} \rightarrow 1^-$, the expression

$$\Xi(\hat{z}) \sim \sum_{\{\pi_1\}} \sum_{\lambda} \rho_h \lambda^{(1-h/2)(\gamma_0-2)-1} = \sum_{\{\pi_1\}} \rho_h \zeta[1 - (1 - h/2)(\gamma_0 - 2)]$$

where $\zeta(\dots)$ denotes Riemann's zeta function. This expression is valid as long as $(1 - h/2)(\gamma_0 - 2) < 0$ for all homotopy types sampled by $\Xi(\hat{z})$. If $(1 - h/2)(\gamma_0 - 2) > 0$,

then it follows that $\Xi(\hat{z})$ behaves critically as

$$\Xi(\hat{z}) \sim \sum_{\{\pi_1\}} \rho_h \frac{\Gamma[(\gamma_0 - 2)(1 - h/2)]}{[c - (1/2)\beta d(\Omega)]^{(\gamma_0 - 2)(1 - h/2)}} \quad (10)$$

when $[c - (1/2)\beta d(\Omega)] \rightarrow 0^+$, (notice that in the above expression $\Gamma(\dots)$ denotes the Euler Gamma function, and that the sum is restricted to those homotopy types for which the above positivity condition holds true).

The analysis of these results is particularly simple and geometrically expressive if we assume that $(\gamma_0 - 2) < 0$, the most general case can be then obtained with obvious modifications.

Under such hypothesis it immediately follows that if all homotopy types sampled in $\mathcal{R}(n, r, D, V)$ by $\Xi(\hat{z})$ have trivial fundamental group, *i.e.*, if $h[\pi_1] = 0$, then as $\hat{z} \rightarrow 1^-$, $\Xi(\hat{z})$ converges to $\sum_{\{\pi_1\}} \zeta[1 - (\gamma_0 - 2)]$. We are sampling simply-connected manifolds, and we have a pure phase. We still have regular behavior if the fundamental group has a non-trivial generator, *i.e.*, if $h[\pi_1] = 1$. But as soon as one of the homotopy type sampled by $\Xi(\hat{z})$ has two non trivial generators, *i.e.*, as $h[\pi_1] = 2$ for some homotopy type, then as $\hat{z} \rightarrow 1^-$ we get a phase transition associated with the simple pole of the corresponding $\zeta[1 - (1 - h/2)(\gamma_0 - 2)]$. Roughly speaking we can interpret this phase transition as the attachment of a handle to the two-skeletons of the manifolds in the homotopy types being sampled. When the number of generators of the fundamental group further increases, a correspondingly increasing number of distinct types of non-simply connected manifolds contributes to the grand-partition function and, as $\hat{z} \rightarrow 1^-$, we move from one to another as if we were attaching the appropriate handles. In this case $\Xi(\hat{z})$ behaves critically as (10).

This criticality stems from the fact that the subleading term $\lambda^{(1 - \frac{h[\pi(M^*)]}{2})(\gamma_0 - 2) - 1}$, in the asymptotics of the counting function B_λ , dominates over the other terms in the statistical sum when we reach the critical temperature.

It is possible to understand this behavior of the grand partition function on more geometrical grounds by estimating, for m very large, the relative mean-square fluctuation in the filling function $N_{(m)}$. An elementary computation provides

$$\frac{\langle (\Delta N_{(m)})^2 \rangle}{\langle N_{(m)} \rangle^2} = \frac{(\sum_{\lambda}^{\sup N_{(m)}} \hat{z}^{\lambda} a_{\lambda})(\sum_{\lambda}^{\sup N_{(m)}} \hat{z}^{\lambda} \lambda^2 a_{\lambda})}{(\sum_{\lambda}^{\sup N_{(m)}} \hat{z}^{\lambda} \lambda a_{\lambda})^2} - 1 \quad (11)$$

where $\langle \dots \rangle$ denotes the average value with respect to the statistical sum $\Xi(m, \hat{z})$, and $\langle (\Delta N_{(m)})^2 \rangle \equiv \langle N_{(m)}^2 \rangle - \langle N_{(m)} \rangle^2$. At the critical temperature, and for $\hat{z} \rightarrow 1^-$, the leading asymptotic behavior of $\langle (\Delta N_{(m)})^2 \rangle / \langle N_{(m)} \rangle^2$, as $m \rightarrow \infty$, is provided by, (again by standard Tauberian theory)

$$\frac{\langle (\Delta N_{(m)})^2 \rangle}{\langle N_{(m)} \rangle^2} = \frac{1}{(1/2)(\gamma_0 - 2)(2 - h[\pi_1(M^*)])} \quad (12)$$

where $h[\pi_1(M^*)] > 2$. Thus, not surprisingly, at the critical regime, the relative root-mean-square fluctuation in the geodesic balls density is not negligible, and we encounter unusually large fluctuations in the filling functions of geodesic balls coverings of manifolds d_G -near M^* . According to (12) such fluctuations are smaller the larger is $h[\pi_1(M^*)]$. This does not surprise, since the larger is the topological complexity of M^* the less *rigid* is the response of M^* to a deformation towards a nearby, (in the Gromov sense), topologically distinct manifold.

4 Connection with 2D-gravity

In order to discuss the connection between the theory presented here and two-dimensional quantum gravity models based on random triangulations of surfaces, let us examine more explicitly the physical meaning of the parameters determining the critical behavior of $\Xi(\hat{z})$. To this end, it is convenient to rewrite Ξ as

$$\Xi(\hat{z}) \sim \rho_h \Gamma[(\gamma_0 - 2)(1 - \frac{h}{2})] \left[\frac{(\beta)^{(2-\gamma_0)}}{[(c/\beta - (1/2)d(\Omega))^{(\gamma_0-2)}]} \right]^{1 - \frac{h[\pi_1(M)]}{2}} \quad (13)$$

This rewriting shows that $\Xi(\hat{z})$, (summed over the finite number of distinct homotopy types realized by manifolds in $\mathcal{R}(n, r, D, V)$), formally has the same structure of the partition function of two-dimensional quantum gravity if we identify $(\beta)^{(2-\gamma_0)}$ with a bare string coupling constant, and c/β , (which, up to a minus sign, is the chemical potential for Ξ), with a bare cosmological constant.

This identification is a consequence of the fact that through $\Xi(\hat{z})$ we are sampling manifold out of $\mathcal{R}(n, r, D, V)$ according to their fundamental group, and this latter is probed by the two-skeleton of the geodesic balls coverings. *As a matter of fact, such two-skeletons are statistically weighted by Ξ through their invariants $N_{(m)}$ and $\Phi_{(m)}$ in the same way as in two-dimensional quantum gravity we weight random triangulation of surfaces.* This explains the similarity between the two theories. In this connection, it is also interesting to remark that, as for two-dimensional gravity, Ξ appears to satisfy a scaling relation: Ξ depends on the ratio

$$\frac{(\beta)^{(2-\gamma_0)}}{[(c/\beta - (1/2)d(\Omega))^{(\gamma_0-2)}]} \quad (14)$$

this remark suggests the question of the existence of a double scaling limit for $\Xi(\hat{z})$ analogous to the double scaling limit in two dimensional quantum gravity which is at the origin of the progresses in such theory, and of the possible relation between such kind of limit and the continuum limit dealt with here.

As is known, the double scaling limit in two-dimensional quantum gravity is taken, as the cosmological constant approaches its critical value, by letting the bare string coupling going to zero while keeping fixed a ratio similar in structure to (14).

It is easily checked that such limit corresponds here to having the ratio (14) finite when, (we keep on assuming that $(\gamma_0 - 2) < 0$), the chemical potential c/β is finite and equal to its critical value $(1/2)d(\Omega)$ and $\beta \rightarrow \infty$. The geometrical meaning of this limiting procedure readily follows from the expression of the critical inverse temperature. This latter diverges if $d_\lambda \rightarrow d(\Omega_{(m)})$ as $m \rightarrow \infty$, namely if we sample manifolds in $\mathcal{R}(n, r, D, V)$ with geodesic balls coverings which are close to being the optimal, coverings, generated by a collection of close packed geodesic balls disposed in a regular array. Roughly speaking, this implies that the limit distribution resulting from this particular limit procedure samples out of $\mathcal{R}(n, r, D, V)$ manifolds which are near constant curvature manifolds M^* , (the given value of the chemical potential fixes the average volume of the manifolds sampled). Double scaling for Ξ thus appears as a particular subcase, (the zero temperature limit), of the possible critical regimes that can be associated with the large m -behavior, on $\mathcal{R}(n, r, D, V)$, of the statistical sum $\Xi(m, \hat{z})$.

4.1 Some concluding remarks

Let us point out some of the problems we face in trying to correlate the theory described by Ξ to anything which could be reasonably called (euclidean) quantum gravity.

A first problem obviously concerns the choice of Gromov's space of bounded geometries as the arena for any quantum theory of gravity. Leaving aside the delicate issue of the Euclidean or Lorentzian signature, the natural objection which comes about Gromov's space concerns the restrictions on curvature, diameter, and volume which appear unnatural in relativity. However, this is not a real issue, for we can remove such bounds by a limit procedure. To be more explicit, we can consider a nested sequence of spaces of bounded geometries

$$\dots \subset \mathcal{R}(n, r_i, D_i, V_i) \subset \mathcal{R}(n, r_{i+1}, D_{i+1}, V_{i+1}) \subset \dots \quad (15)$$

where $r_i \rightarrow \infty$, $D_i \rightarrow \infty$, and $V_i \rightarrow 0$. In this way we may formally recover the space of all riemannian structures (completed under Gromov-Hausdorff convergence) as a limit of spaces of bounded geometries. *Manifolds* in such space may have arbitrarily large curvatures, arbitrarily small and/or large volumes and diameters, and we can consider the behavior of the associated sequence of partition functions Ξ_{n, r_i, D_i, V_i} . The limiting behavior of such sequence appears quite non-trivial, if, for instance we consider each Ξ_i at its critical point, where topology dominates. Simplyfing a little bit, we can say that the philosophy underlying the use of the compact Gromov spaces is simply that such spaces allow for a clear mathematical control of the *cut-off parameters* which determine the behavior, in the continuum limit, of symplcial approximations to manifolds.

A more delicate issue with Ξ concerns the *regularity* of the manifolds sampled out in $\mathcal{R}(n, r, D, V)$ according to Ξ . More explicitly: when are the (homology) manifolds, dominating in Ξ , *smooth* differentiable manifolds? This problem is already present in

two-dimensional quantum gravity, where it takes the form of the non-vanishing of the effective string coupling constant at the critical point. This yields for a continuum limit where the dominating *surface* configurations are those of *crumpled* objects where diffeomorphism invariance is completely lost. Here, the situation is not very different, in place of crumpled surfaces we, generally speaking, have to deal with homology manifolds (in the physically interesting dimensions three and four), or with topological manifolds as soon as the dimension equals five. According to the results of the previous paragraphs we have some control on what happens at the critical point to such homology manifolds, but we have no explicit results which enforce the onset of diffeomorphism invariance. A rather shy indication in such direction comes from the geometrical meaning of the double scaling limit which can be implemented for Ξ , but admittedly this is not much.

References

- 1 E.Brézin and V.Kazakov, Phys. Lett. **B236**,144 (1990).
- 2 M.Douglas and S.Shenker, Nucl. Phys. **B235**, 635 (1990).
- 3 D.J.Gross and A.A.Migdal, Phys. Rev. Lett. **64**, 127 (1990); Nucl. Phys. **B340**, 333 (1990).
- 4 M.Carfora and A.Marzuoli, *Bounded geometries and topology fluctuations in lattice quantum gravity* to appear in Class.Quant. Grav. (1991); see also *Critical phenomena for Riemannian geometries* Preprint FNT/T 90/07, (Univ. Pavia).
A more detailed mathematical presentation can be found in
M.Carfora and A.Marzuoli, *Finiteness theorems in Riemannian geometry and lattice quantum gravity*, to appear in Contemporary Mathematics (Proceedings of the AMS research meeting *Mathematical Aspects of Classical Field Theory*, Seattle, 1991).
- 5 M.Carfora and A.Marzuoli, Phys. Rev. Lett. **62**, 1339 (1989).
- 6 M.Gromov, *Structures métriques pour les variétés Riemanniennes* (Conception Edition Diffusion Information Communication Nathan, Paris, 1981).
See also S.Gallot, D.Hulin, J.Lafontaine *Riemannian Geometry*, (Springer Verlag, New York,1987).
For the convenience of the reader we recall that the diameter of a riemannian manifold M is defined as $\sup_{(p,q) \in M \times M} d(p,q)$ where $d(\cdot, \cdot)$ denotes the distance function of M .
- 7 K.Grove and P.V.Petersen, Annals of Math. **128**, 195 (1988).

- 8 P.Erdos, P.Kelly, Amer.Math.Monthly **70** 1074 (1963); see also chap.10 of F.Harary (ed.) *A Seminar on Graph Theory* (Holt, Rinehart and Winston, Inc.,New York, 1967).
- 9 D.Ruelle,*Statistical Mechanics: rigorous results* (W.A.Benjamin, New York, 1974).
- 10 D.Bessin, C.Itzykson and J.B.Zuber, Adv. Appl. Math. **1**,109(1980).
- 11 F.Harary and E.Palmer, *Graphical Enumeration* (Academic, New York, 1973).
- 12 K.Grove, P.V.Petersen and J.Y.Wu, Bull. Am. Math. Soc. **20**, 181 (1989).
- 13 J.Ambjorn, *Random surfaces: a non-perturbative regularization of strings in Probabilistic Methods in Quantum Field Theory and Quantum Gravity* P.H.Damgaard *et al.* eds. (Plenum Press, New York, 1990).
- 14 K.Grove, P.V.Petersen and J.Y.Wu, Bull.Am.Math.Soc. **20**, 181 (1989); Invent.Math. **99**, 205 (1990) (and its Erratum).
- 15 It is interesting to note that the structure of $\mathcal{R}(n, r, D, V)$ is quite sensible to the dimension n , in the sense that for $n \geq 5$ one can actually count (at least in principle), the different types of topologies and the different types of differentiable structures realized in each homotopy class of manifolds sampled by $\Xi(m, z)$, since the number of such structures can be shown to be finite. In dimension three and four this counting is no longer possible. Some remarks concerning the phase structure of high-dimensional, ($n \geq 5$), lattice quantum gravity theories will be discussed elsewhere.